

VAYSMAN, M.L.; TROYNO, V.P.; PERTSEL', V.M.

Use of ultrasound in the control of scale formation in evaporators. Sakh.prom. 34 no.1:36-39 Ja '60.

(MIRA 13:5)

1. TSentral'nyy nauchno-issledovatel'skiy institut sakharnoy promyshlennosti (for Vaysman, Troyno). 2. 2-y Petrovskiy sakharnyy zavod (for Pertsel').

(Sugar manufacture)

(Ultrasonic waves--Industrial applications)

KHONIG, P.[Honig, Pieter], red.; GOLOVNYAK, Yu.D., inzh.[translator];
MAKSIMOVA, N.A., inzh. [translator]; ZHIZHINA, R.G., inzh.
[translator]; Prinimali uchastiye: TROYNO, V.P. [translator];
GOROKH, V.N.[translator]; BENIN, G.S., kand. tekhn. nauk, red.;
VOYKOVA, A.A., red.; KISINA, Ye.I., tekhn. red.

[Principles of sugar technology]Printsiipy tekhnologii sakhara.
Pod red. G.S.Benina. Moskva, Pishchepromizdat, 1961. 615 p.
Translated from the English. (MIRA 15:12)
(Sugar manufacture)

TROYNO, V.P.

Determining the velocity profile in the downtake tube of the
massecuite vacuum apparatus. Izv.vys.ucheb.zav.; pishch.tekh.
no.3:116-122 '62. (MIRA 15:7)

1. Kiyevskiy tekhnologicheskoy institut pishchevoy promyshlennosti,
kafedra teploenergetiki.
(Sugar manufacture) (Vacuum apparatus—Fluid dynamics)

TROYNO, V.P.; POPOV, V.D.

Effect of the liquid level and circulation rate on heat
exchange in the boiling of massecuite. Trudy KTIPP no.25:
89-98 '62. (MIRA 16:5)
(Vacuum apparatus) (Heat—Transmission) (Sugar manufacture)

TROYNO, V. P.; VAYSMAN, M. L.

Temperature and height of the boiling point of massecuite.
Izv.vys.ucheb.zavl; pishch.tekh.no. 2:128-130 '64. (MIRA 17:5)

POPOV, V.D., doktor tekhn. nauk; TROYNO, V.P., kand. tekhn. nauk

Hydraulic resistance in the flow of sugar massecuite. Pishch.
prom. no.1:122-130 '65. (MIRA 12:11)

PROYNO, V.F., kand. tekhn. nauk; POPOV, V.L., doktor tekhn. nauk

Hydrodynamic design of sugar refining vacuum apparatus for
periodical and continuous action. Pishch. prom. no. 10/197-115
'65. (MIRA 18:1)

1. Kiyevskiy tekhnologicheskii institut pishchevoy promyshlen-
nosti.

TROZCENKO, A. T.

"Syntheses en partant des o- et p-dioxydiphenyles. Memoire III". Vorozcov, N. N. (junior).
Troscenko, A. T. (p. 59)

SO: Journal of General Chemistry
(Zhurnal Obshchei Khimii) 1939, Volume 9, #1

TROZOS, A.

"Preparing Trucks for Winter." p. 301, (MOTORYZACJA, Vol. 8, No. 11,
Nov. 1953. Warszawa, Poland.)

SO: Monthly List of East European Accessions, (EEAL), LC,
Vol. 3, No. 12, Dec. 1954, Uncl.

TROZYAN, R.Ye.

Calculation of the water hammer in pipelines. Izv. AN Arm.
SSR. Ser. tekhn. nauk 16 no.4:70-72 '63. (MIRA 16:10)

YUGOSLAVIA/Chemical Technology. Chemical Products and Their Application. Electrochemical Industries. Electroplating Galvanic Cells

Abs Jour : Ref Zhur - Khimiya, 1958, No 22, 74616

Author : Brchich B., Moyech B., Trpevske B.

Inst : Not Given

Title : Anodic Oxidation of Ferrochrome

Orig Pub : Glasnik Khim. drushtva, 1957, 22, No 4, 233-243

Abstract : Anodic oxidation of ferrochrome was investigated employing solutions of NaCl, Na₂CrO₄, and H₂CrO₄ of varying concentrations. An increase in BT_a was observed when concentration of the electrolyte (NaCl) and D_a were reduced. The optimum conditions of oxidation ($BT_a \sim 73\%$) were as follows: NaCl concentration of 0.05n, $D_a = 0.5$ a/ m², mixing with air. At those conditions the Cr³⁺ content was reduced to 8%. The Fe:Cr ratio in the electrolyte differs from that on the anode. With the decreased concentration of the electrolyte, the Fe concentration in the solution decreases also. At D_a 1a/ m² and while mixing with air the Fe:Cr ratio in the

Card : 1/3

YUGOSLAVIA/Chemical Technology. Chemical Products and Their Appli- H-12
cation. Electrochemical Industries. Electroplating Galvanic
Cells

Abs Jour : Ref Zhur - Khimiya, 1958, No 22, 74616

0.05 n NaCl solution and on the anode are equal. At lower values of D_c (0.5 s/cm^2) the electrolyte contains less Fe. At the other D_a values and at the same NaCl concentration; as well as at all the values of D_a and in the 2.7 n NaCl solution, the electrolyte contains more Fe than it is present on the anode. Similar behavior was observed with the 0.025 n H_2CrO_4 solutions in which the Cr^{3+} content was equal to 16.4%. pH of the electrolytes falls rapidly during the first 10-15 amp. hrs. from 7 to 1.5-2.0. In experimenting with the 0.1 n H_2CrO_4 solution it was observed that under certain conditions, value of the electrolyte pH increases and at a pH of approx. 2.0, $\text{Fe}(\text{OH})_3$ precipitates. In the presence of $\text{Cr}_2\text{O}_7^{2-}$ ions electrical charge of the above precipitate changes and Fe is deposited on the anode. Thickness of the formed layer depends on the dispersion of $\text{Fe}(\text{OH})_3$ and determines the degree of resistance thus produced. Such a

Card : 2/3

YUGOSLAVIA/Chemical Technology. Chemical Products and Their Appli- H-12
cation. Electrochemical Industries. Electroplating Galvanic
Cells.

Abs Jour : Ref Zhur - Khimiya, 1958 , No 22, 74616

phenomenon may be avoided by increasing concentration of H_2CrO_4 up to 1 n. Under these conditions pH of the solution remains below 2 for a prolonged time. The described phenomenon was not observed in neutral solutions. In the latter case $Fe(OH)_3$ was found to have high dispersivity and the Cr^{3+} content in such solutions was approx. 15%. At elevated temperatures resistance of the electrolytes decreases, which is particularly advantageous in the initial stages of the process. Consumption of the electric energy in all the cases was found almost identical and comprised 4.7 KW Hrs for 1 kg CrO_3 or 2.4 KW Hrs for 1 kg K_2CrO_4 .

Card : 3/3

TRPENOVSKI, Branko; CUPONA, Gorgi

Finitary associative operations with neutral elements.
Bilten mat fiz Mak no.12:15-24 '61

PAVLOVIC, V.; TRPINAC, P.

Determination of the structure of dextran by the oxidation
of periodate. Vojnosanit pregl 19 no.7/8 J1-Ag '62.

1. Medicinski fakultet, ~~Un~~iverzitet a Beogradu.
Hemijski institut.
(PERIODIC ACIDS) (DEXTRAN)

PAVLOVIC, V.; TRPINAC, P.

Determination of the structure of dextran by the oxidation
of periodate. Vojnosanit. pregl. 19 no.7/8:542-545 JI-Ag '62.

1. Medicinski fakultet u Beogradu, Hemijski institut.
(PERIODIC ACIDS) (DEXTRAN)

S

SULOVIC, Vojin; BUGARSKI, Olga; RCTOVIC, Bozica; TRPINAC, Pavle; SKURINA, Tatjana

Electrophoresis of serum proteins in early and late pregnancy toxemias. Srpski arh. celok. lek. 89 no.12:1435-1442 D '61.

1. Hemijski institut Medicinskog fakulteta Univerziteta u Beogradu
Upravnik: prof. dr Pavle Trpinac Ginekološko-akuserska klinika Medicinskog fakulteta Univerziteta u Beogradu Upravnik: prof. dr Bosiljka Milosevic.

(PREGNANCY TOXEMIAS blood)
(BLOOD PROTEINS impregn)
(ELECTROPHORESIS)

YUGOSLAVIA

A. LAZAROVIC and P. THOMAS, Department of Chemistry, Medical Faculty
(Cetinjski put 131) (Medicinsko-farmaceutski fakultet) University of Belgrade

"Study of Percentage of (1,6) Bonds in PUDT Clinical Dextran by Means
of Periodate Oxidation."

Belgrade, Arhiv za Higijenu, Vol 12, No 4, 1962; pp 221-223;

Abstract [French summary modified]: Study of Yugoslav-made dextran
(Ing. A. and L. Vavra, Novi Sad) by periodate oxidation method to
determine percentage of alpha (1,6) bonds by periodate consumed to
formic acid formed per mole. Domestic dextran (specimens from 2 batches)
was as good as the Swedish- or Swiss-made product tested as controls.
Structural formula, table; 6 Western references.

1/1

TRPINAC, Pavle, prof. d-r

Clinical laboratory in the current stage of development of our health services. Voj. san. pregl, Beogr. 16 no7-8:583-584 J1-Ag '59.

1. Medicinski fakultet u Beogradu, Hemiski institut.
(LABORATORIES)

TRPINAC, Pavle, prof., dr.

Standardization of clinico-chemical methods. Voj.san.pregl. 18 no.5:
443-444 My '61.

1. Medicinski fakultet u Beogradu, Hemijski institut.

(DIAGNOSIS LABORATORY)

TRPINAC, Pavle Dr.

The collaboration of physicians and pharmacists in the improvement
of public health service. Arh.farm., Beogr. 5 no.1:1-9 Feb. 55.
(PUBLIC HEALTH,

in Yugosl., improvement by collaboration of physicians
& pharmacists (Ser))

TRIAS, M.

SCIENCE

TRIAS, M. Notes on the ecology and zoogeography of the species *Aedes* (C.)
refiki (Diptera, Culicidae). p. 305.

Vol. 13, No. 4, 1958.

Monthly Index of East European Accessions (EEAI: IC, Vol. 7, No. 12, Dec. '58

1 Rpt 11
CZECHOSLOVAKIA / General and Special Zoology. Insects. P
Systematics and Faunistics.

Abs Jour: Ref Zhur-Biol., No 21, 1958, 96359.

Author : Trpis, M.

Inst : Not given.

Title : Preliminary Survey of Dragonflies on Zhitnyy
Island.

Orig Pub: Biologia, 1957, 12, No 6, 433-449.

Abstract: The island of Zhitnyy is located on the Danube
lowland to the east of Bratislava. 46 species of
dragonflies were found on the island. The fauna
of the island dragonflies is basically Central
European with a large admixture of Mediterranean
species. -- From the author's resume.

Card 1/1

TRPI3, M.

"Research on the natural focal points of communicable diseases in Yugoslavia."

p. 149 (Biologia, Vol. 13, no. 2, 1958, Praha, Czechoslovakia)

Monthly Index of East European Accessions (EEAI) LC, Vol. 7, no. 7,
September 1958

TRPIS, M.

Experiences in fighting mosquitoes in southwestern Slovakia. p.27.
(BIOLOGICKE PRACE, Vol. 2, no. 6, 1957, Bratislava, Czechoslovakia.)

SO: Monthly List of East European Accessions (EEAL) LC, Vol. 6, no. 12, December 1957. Incl

TRPIS, Milan

Determination of the number of mosquitoes in eastern Slovakia.
Biologia (Bratisl.) 19 no.11:243-248 '64

1. Abteilung für Zoologie des Biologischen Institutes der
Slowakischen Akademie der Wissenschaften in Bratislava.

as Disease Vectors.

Abs Jour: Ref Zhur-Biol., No 9, 1958, 38624.

Author : Trpis, M.

Inst : ~~Not given.~~

Title : Experimental Mosquito Control in Southwestern Slovakia.

Orig Pub: Biol. prace, 1956, 2, No 6, 27-46.

Abstract: Control of mosquitoes *Aedes vexans*, bred in flood waters of Zhitny Island, was conducted by treating that locality with dinocide (a preparation containing 5% DDT) at the rate of 0.1 - 1.0 gm/m², from airplanes. Altogether 14,848 hectares were treated. In the summer of 1954 the number of mosquitoes in flooded woods before treatment consisted of nearly 50,000 per km (sic). After treatment

Card 1/2

CZECHOSLOVAKIA / Zooparasitology - Mites and Insects G-3
as Disease Vectors.

Abs Jour: Ref Zhur-Biol., No 9, 1958, 38624.

Abstract: the number diminished by 98.47% and remained at
this level for 5-7 days, after which it gradually
increased.

Card 2/2

L 00057-66 EMT(1)/T/EMA(b)-2 BW/JK

ACCESSION NR: AP5023866

CZ/0049/04/000/011/0843/0848

AUTHOR: Trpish, Milan (Trpish, Milan) (Graduate biologist, Candidate of sciences) (Bratislava)

TITLE: Areas of occurrence of mosquitoes in Eastern Slovakia shown with frequency of quantitative distribution

SOURCE: Biologia, no. 11, 1964, 843-848

TOPIC TAGS: parasitology, animal parasite, entomology

ABSTRACT: The distribution of mosquitoes in Eastern Slovakia varies according to conditions in individual regions. In the low-lying areas near the river Tisa, yearly flooding of woodlands occurs. After research lasting 3 years the authors divided Eastern Slovakia into 4 regions: 1. Region with a prevalence of mosquitoes every year. 2. Region with irregular prevalence of mosquitoes. 3. Region of high incidence of mosquitoes. 4. Region with low incidence of mosquitoes. A map showing the 4 regions is presented. This map should help in the fight against the mosquitoes. / Orig. art. has: 1 figure.

Card 1/2

L 00057-66

ACCESSION NR: AP5023866

3

ASSOCIATION: Abteilung fur Zoologie des Biologischen Institutes der Slowakischen Akademie der Wissenschaften, Bratislava (Department of Zoology, Institute of Biology, Slovak Academy of Sciences) 4055

SUBMITTED: 08 Jun 64

ENCL: 00

SUB CODE: LS

NR REF SOV: 000

OTHER: 003

JPRS

mlh
Card 2/2

"APPROVED FOR RELEASE: 03/14/2001

CIA-RDP86-00513R001756810008-5

APPROVED FOR RELEASE: 03/14/2001

CIA-RDP86-00513R001756810008-5"

TRPIS, Milan

On some parasitological problems in Rumania. Biologia 17 no.11:849-854 '62.

1. CSAV, Biologicky ustav Slovenskej akademie vied, Oddelenie zoologie v Bratislave.

(PARASITIC DISEASES)

ROMANIA

Milan TREIS, Department of Zoology, Biological Institute of the Slovak Academy of Sciences, Czechoslovak Academy of Sciences (oddelenie zoologie, Biologický ústav Slovenskej akadémie vied, Československá akadémia vied) Bratislava.

"Some Parasitological Problems in Romania."

Bratislava, Biologia, Vol 17, No 11, 1962: 11 849-854.

Abstract: Author spent 3 weeks in Romania (Aug. 1961) as guest of the Cantacuzino Institute to study malaria and mosquitoes in the marshes; his article briefly reviews the history of the Institute and some of its general activities, but concentrates on the problems of mosquitoes and malaria. Massive planned applications of insecticides have greatly decreased the incidence of malaria during the recent years despite the greater difficulty in Romania due to topographical factors. Four photographs.

1/1

TRPIS, Milan

Entomologic Days. Biologia 15 no.12:948-949 '60. (EEAI 10:8)
(CZECHOSLOVAKIA—ENTOMOLOGY)

TRPIS, Milan

New informations on a method for the study on activities of mosquitoes.
Biologia 17 no.2:123-129 '62.

1. CSAV - Biologisches Institut der Slowakischen Akademie der Wissenschaften, Abteilung fur Zoologie, Bratislava.

(MOSQUITOES)

TRPIS, Milan

1st detection of *Theobaldia* (A.) *longiareolata* Macq. 1838 (Diptera, Culicidae) in Czechoslovakia. *Biologia* 17 no.3:213-215 '62.

1. CSAV - Biologicky ustav Slovenskej akademie vied, Oddelenie zoologie v Bratislave.

(DIPTERA)

TRPTS, M.; TOVORNIKOVA, D.

Faunistic, ecologic, and zoogeographic remarks on mosquitos in Slovenia, Yugoslavia. In German. p. 721

BIOLOGIA. (Slovenska akademia vied) Bratislava, Czechoslovakia, Vol. 13, no. 10 1958

Monthly List of East European Acquisitions (MEAI), LC, Vol. 8, no. 11, Nov. 1959
Uncl.

Trpis, M.; Korbel, L.

Report on the 2d session of the Czechoslovak Entomologic Society in Slovakia at the Slovak Academy of Sciences. p. 550.

BIOLOGIA, Bratislava, Czechoslovakia, Vol. 14, no. 7, 1959

Monthly List of East European Accessions (EEAI) LC, Vol. no. ⁸10, 1959 -Oct.
Uncl.

TRPIS, M.

Trpis, M. Mosquitos in the High Tatra (Diptera, Culicidae). p.221.

Vol. 10, no.2, 1955 BIOLOGIA Bratislava, Czechoslovakia

SO: Monthly List of East European Accessions, (EEAL), LC, Vol. 5, No.2
February, 1956

TRPIS, M.

Mosquitoes in the Vah River valley (Diptera, Culcidae). p. 507.

BIOLORGIA. (Slovenska akademia vied) Bratislava (CZECHOSLOVAKIA)

Vol, 10, No. 4, 1955.

SOURCE: East European Accessions List (EEAL) Library
of Congress. Vol. 5, No. 1, January, 1956.

TRPIS, Milan

Investigations on mosquitoes in high Taters (Diptera, Culicidas).
Biologia, Bratisl. 10 no.2:231-236 '55.

1. Faunisticke laboratorium Slovenskij akademie vied a Zoologicky
ustav University Komenskeho v Bratislave.

(MOSQUITOES,

distribution in Czech. mountain region)

TRPIS, M.

A preliminary survey of dragonflies (Odonata) on Litny Ustrov.

P. 433, (Biologia) Vol. 12, no. 6, 1957, Praha, Czechoslovakia.

SO: Monthly Index of East European Acessions (EEAI) Vol. 6, No. 11 November 1957

TRPIS, Milan

Mosquitoes (Diptera, Culicidae) in the Bratislava region. Biologia,
Bratisl. 9 no.4:412-424 1954.

1. Zoologicky ustav SU v Bratislave.
(MOSQUITOES,
Culex, in Czech.)

TRPIS, Milan

Activity and seasonal dynamics of flies on the locations of their
hiding-places in the vegetation of the Danube valley forests.
Biologia 17 no.4:263-282 '62.

1. CSAV, Biologicky ustav Slovenskej akademie vied, Oddelenie
zoologie v Bratislave.
(DIPTERA) (HOUSEFLIES)

TRPIS, Milan

Some new information on the construction of light traps for insects. Biologia (Bratisl.) 20 no.12:901-907 '65.

1. Oddelenie ekologickej fyziologie hmyzu Ustavu biologie krajiny Slovenskej akademie vied v Bratislave.

ca

2,4,6-Trichloro-3-amino-1-methylbenzene and some
of its derivatives. E. Burel and M. Trpilovski. *Caspis*
Czechoslov. Liharska 15, 179-~~180~~(1955). By the action
of Cl on m-acetotoluene in glacial AcOH there was ob-
tained 2,4,6-trichloro-3-acetamino-1-methylbenzene, m.
192°. The sapon of this product gave 2,4,6-trichloro-
m-toluidine, m. 85°. By the nitration of 2,4,6-Cl₃C₆H₃Me
there was obtained 2,4,6-trichloro-3-nitrotoluene, m. 50°
and by the nitration of 2,3,4,6-Cl₄C₆HMe the resulting
product was 2,3,4,6-tetrachloro-5-nitrotoluene, m. 148-
50°.

V. D. Karpenko

BC

2:4:6-Trichloro-*m*-toluidine and some derivatives. E. Bunzl and M. Taribovská (Časopis českých věd., 1935, 15, 179—185; Chem. Zentr., 1936, 1, 1209).—Chlorination of acet-*m*-toluidide in AcOH affords 2:4:6-trichloroacet-*m*-toluidide, m.p. 102°, hydrolyzed (NaOH) to 2:4:6-trichloro-*m*-toluidine, m.p. 85° (lit. m.p. 218°, and Ac derivative, m.p. 81—83°). 2:4:6-Trichloro-*m*-toluidine, m.p. 38°, and 2:3:4:6-tetrachloro-*m*-toluidine, m.p. 91.5—92°, toluene are prepared from the appropriate amines by the diazo reaction; on nitration they yield 2:4:6-trichloro-3-nitrotoluene, m.p. 50°, and 2:3:4:6-tetrachloro-3-nitrotoluene, m.p. 148—150°, respectively. 2:4:6-Trichloro-3-bromo-*m*-toluidine, m.p. 85°, and 2-iodo-*m*-toluidine, m.p. 63°, are obtained from the appropriate diazonium salts, Cu-bromide, and KBr or KI. H. N. R.

TRPKOS, L.

TRPKOS, L. Founding of cylinders for air-cooled mo tors. p. 375

Vol. 10, no. 12, 1956, June
SVET MOTORU
TECHNOLOGY
Praha, Czechoslovakia

So: East European Accession Vol. 6, no. 2, 1957

TRPKOS, Ladislav, inz.

Trail car tractor Praga S5T-2-TN. Siln doprava 11 no.11:
10-12 N '63.

1. Vyvoj automobilu, Automobilove zavody, Letnany.

TRFKOVIC, Miodrag

"Burnishing of iron and steel"

SO: TEHNIKA No 7, Year X, - 1955¹

TRPLOVIC, A.

TRPLOVIC, A. The most suitable method for testing materials without breaking them. p. 14

Vol. 12, no. 12, Dec. 1956

ZELEZNICE

TECHNOLOGY

Beograd

So: East European Accession, Vol. 6, no. 3, March 1957

TRFUTEC, V.

Yugoslavia (430)

Technology-Periodicals

Flaw formations in aluminum alloys. p. 294. TEHNICKI PREGLED. (Croatia. Uprava za unapredenje proizvodnje pri privednom savjetu) Zagreb. (Bimonthly technical journal issued by the Production Improvement Administration of the Economic Council) No. 6, 1951.

East European Accessions List. Library of Congress Vol. 2, No. 6, June 1953. Unclassified.

TRSINSKI, M.

See Drezancic, I.

TRSOHIN, Ye. I.; AFONIN, G. G.

Temperatures of the formation of pyrrhotites from certain complex
metal deposits in Transbaikalia. Geokhimiya no. 11: 2199-2200 N 164.
(MIRA 18:8)

1. Institut geokhimiya Sibirskogo otdeleniya AN SSSR, Irkutsk.

4

PROCESSES AND PROPERTIES INDEX

PAPIR-ES NYOMDATECHNIKA -- PAPER AND PRINTING
Vol. 2 -- 1950
No. 9, Sept.

34

by *Erstgonyak* 311215
Raising basic wages and adjusting
norms p. 21

ASAC SEA METALLURGICAL LITERATURE CLASSIFICATION

REPORT NO. 101 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100

CLASSIFICATION 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100

CZECHOSLOVAKIA, Cultivated Plants - Fruits. Berries.

M.

Abstr Jour : Ref Zhur - Biol., No 10, 1958, 44340

Author : Trestenjak, Milko

Inst : -

Title : Grape Selection in 1956.

Orig Pub : Sadjar., vinar., vrbnar., 1957, 44, No 4, 110-112

Abstract : The Institute of Horticulture and Viticulture in Maribor has been conducting large scale experiments since 1956 on the selection of grape on 15 sections in different regions of Slovenia. In Maribor the experiments were started in 1947 and by 1956 the number of selected bushes of 30 varieties reached 120000. -- Ye.A. Parshina

Card 1/1

- 174 -

TRSTENJAK, Miso

Present state of the physics of the electromagnetic field. Elektr vest
27 no.11/12:428-429 N-D '59. (EEAI 10:1)
(Electromagnetism) (Quantum theory)

STICKY, Petr; TRTIK, Josef

Casting of worm wheels from aluminum bronze. Slevarenstvi
10 no.7:266-267 J1 '62.

1. Choteborske kovodelne zavody, Nove Ransko.

JELMANOV, Ivan, inz.; PISEK, Jaroslav, inz.; TRSEK, Miroslav, inz.

Boring with local circulation at the borehole bottom. Geol
pruzkum 7 no.2:51-52 F '65.

1. Jachymovske doly, Geologicky pruzkum National Enterprise,
Pribram.

CA

7

Determination of traces of iodine. Josef Trtlik.
Chem. Listy 38, 128-31 (1944); cf. *C.A.* 34, 1000. To
improve the accuracy of the detn. of I by mercurimetric
titration with dithizone as indicator, T. suggests compar-
ison of the coloration of the titrated soln. with a blank
contg. a trace of $\text{Hg}(\text{NO}_3)_2$. With 0.004 and 0.002 N
solns. 2 mg. of I were detd. with 0.1-1% accuracy.
Milos Hudlicky

BC

17-1

Application of diphenylcarbazide and diphenylcarbazone to mercurimetric microtitration. J. V. Donsky and J. Terfink (Chem. Listy, 1933, 27, 385-388).—Diphenylcarbazide and -carbazone give intense violet colorations in presence of Hg^{++} . Accurate results are obtained using these indicators in titration of Cl^- by $0.01N-Hg(NO_3)_2$; Cu^{++} , Cd^{++} , Co^{++} , Fe^{++} , Ca^{++} , and Ba^{++} do not interfere. The solution titrated should contain a small excess of HNO_3 .

R. T.

ASSOCIATED METALLURGICAL LITERATURE CLASSIFICATION

SECTION ONE										SECTION TWO																																																																																									
SUBSECTION ONE										SUBSECTION TWO																																																																																									
SUBSECTION ONE										SUBSECTION TWO																																																																																									
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100

[illegible]

MATERIALS		PROCESSING AND PROPERTIES INDEX	
GROUP	ELEMENT	GROUP	PROPERTY
1	Al	1	Al
2	Ca	2	Ca
3	Fe	3	Fe
4	Co	4	Co
5	Ni	5	Ni
6	Cu	6	Cu
7	Zn	7	Zn
8	Mn	8	Mn
9	Mg	9	Mg
10	Na	10	Na
11	K	11	K
12	Li	12	Li
13	Be	13	Be
14	B	14	B
15	C	15	C
16	N	16	N
17	O	17	O
18	F	18	F
19	Cl	19	Cl
20	S	20	S
21	P	21	P
22	As	22	As
23	Sb	23	Sb
24	Bi	24	Bi
25	Pb	25	Pb
26	Sn	26	Sn
27	Hg	27	Hg
28	Ag	28	Ag
29	Au	29	Au
30	Pt	30	Pt
31	Ir	31	Ir
32	Rh	32	Rh
33	Os	33	Os
34	Re	34	Re
35	Mo	35	Mo
36	Ta	36	Ta
37	Nb	37	Nb
38	W	38	W
39	Cr	39	Cr
40	Mn	40	Mn
41	Fe	41	Fe
42	Co	42	Co
43	Ni	43	Ni
44	Cu	44	Cu
45	Zn	45	Zn
46	Ga	46	Ga
47	Ge	47	Ge
48	As	48	As
49	Se	49	Se
50	Br	50	Br
51	I	51	I
52	At	52	At
53	H	53	H
54	He	54	He
55	Li	55	Li
56	Be	56	Be
57	B	57	B
58	C	58	C
59	N	59	N
60	O	60	O
61	F	61	F
62	Ne	62	Ne
63	Na	63	Na
64	Mg	64	Mg
65	Al	65	Al
66	Si	66	Si
67	P	67	P
68	S	68	S
69	Cl	69	Cl
70	Ar	70	Ar
71	K	71	K
72	Ca	72	Ca
73	Sc	73	Sc
74	Ti	74	Ti
75	V	75	V
76	Cr	76	Cr
77	Mn	77	Mn
78	Fe	78	Fe
79	Co	79	Co
80	Ni	80	Ni
81	Cu	81	Cu
82	Zn	82	Zn
83	Ga	83	Ga
84	Ge	84	Ge
85	As	85	As
86	Se	86	Se
87	Br	87	Br
88	Kr	88	Kr
89	Rb	89	Rb
90	Sr	90	Sr
91	Y	91	Y
92	Zr	92	Zr
93	Nb	93	Nb
94	Mo	94	Mo
95	Tc	95	Tc
96	Ru	96	Ru
97	Rh	97	Rh
98	Pd	98	Pd
99	Ag	99	Ag
100	Cd	100	Cd
101	In	101	In
102	Sn	102	Sn
103	Pb	103	Pb
104	Bi	104	Bi
105	Po	105	Po
106	At	106	At
107	Hg	107	Hg
108	Tl	108	Tl
109	Pb	109	Pb
110	Bi	110	Bi
111	Po	111	Po
112	At	112	At
113	Hg	113	Hg
114	Tl	114	Tl
115	Pb	115	Pb
116	Bi	116	Bi
117	Po	117	Po
118	At	118	At
119	Hg	119	Hg
120	Tl	120	Tl
121	Pb	121	Pb
122	Bi	122	Bi
123	Po	123	Po
124	At	124	At
125	Hg	125	Hg
126	Tl	126	Tl
127	Pb	127	Pb
128	Bi	128	Bi
129	Po	129	Po
130	At	130	At
131	Hg	131	Hg
132	Tl	132	Tl
133	Pb	133	Pb
134	Bi	134	Bi
135	Po	135	Po
136	At	136	At
137	Hg	137	Hg
138	Tl	138	Tl
139	Pb	139	Pb
140	Bi	140	Bi
141	Po	141	Po
142	At	142	At
143	Hg	143	Hg
144	Tl	144	Tl
145	Pb	145	Pb
146	Bi	146	Bi
147	Po	147	Po
148	At	148	At
149	Hg	149	Hg
150	Tl	150	Tl
151	Pb	151	Pb
152	Bi	152	Bi
153	Po	153	Po
154	At	154	At
155	Hg	155	Hg
156	Tl	156	Tl
157	Pb	157	Pb
158	Bi	158	Bi
159	Po	159	Po
160	At	160	At
161	Hg	161	Hg
162	Tl	162	Tl
163	Pb	163	Pb
164	Bi	164	Bi
165	Po	165	Po
166	At	166	At
167	Hg	167	Hg
168	Tl	168	Tl
169	Pb	169	Pb
170	Bi	170	Bi
171	Po	171	Po
172	At	172	At
173	Hg	173	Hg
174	Tl	174	Tl
175	Pb	175	Pb
176	Bi	176	Bi
177	Po	177	Po
178	At	178	At
179	Hg	179	Hg
180	Tl	180	Tl
181	Pb	181	Pb
182	Bi	182	Bi
183	Po	183	Po
184	At	184	At
185	Hg	185	Hg
186	Tl	186	Tl
187	Pb	187	Pb
188	Bi	188	Bi
189	Po	189	Po
190	At	190	At
191	Hg	191	Hg
192	Tl	192	Tl
193	Pb	193	Pb
194	Bi	194	Bi
195	Po	195	Po
196	At	196	At
197	Hg	197	Hg
198	Tl	198	Tl
199	Pb	199	Pb
200	Bi	200	Bi
201	Po	201	Po
202	At	202	At
203	Hg	203	Hg
204	Tl	204	Tl
205	Pb	205	Pb
206	Bi	206	Bi
207	Po	207	Po
208	At	208	At
209	Hg	209	Hg
210	Tl	210	Tl
211	Pb	211	Pb
212	Bi	212	Bi
213	Po	213	Po
214	At	214	At
215	Hg	215	Hg
216	Tl	216	Tl
217	Pb	217	Pb
218	Bi	218	Bi
219	Po	219	Po
220	At	220	At
221	Hg	221	Hg
222	Tl	222	Tl
223	Pb	223	Pb
224	Bi	224	Bi
225	Po	225	Po
226	At	226	At
227	Hg	227	Hg
228	Tl	228	Tl
229	Pb	229	Pb
230	Bi	230	Bi
231	Po	231	Po
232	At	232	At
233	Hg	233	Hg
234	Tl	234	Tl
235	Pb	235	Pb
236	Bi	236	Bi
237	Po	237	Po
238	At	238	At
239	Hg	239	Hg
240	Tl	240	Tl
241	Pb	241	Pb
242	Bi	242	Bi
243	Po	243	Po
244	At	244	At
245	Hg	245	Hg
246	Tl	246	Tl
247	Pb	247	Pb
248	Bi	248	Bi
249	Po	249	Po
250	At	250	At
251	Hg	251	Hg
252	Tl	252	Tl
253	Pb	253	Pb
254	Bi	254	Bi
255	Po	255	Po
256	At	256	At
257	Hg	257	Hg
258	Tl	258	Tl
259	Pb	259	Pb
260	Bi	260	Bi
261	Po	261	Po
262	At	262	At
263	Hg	263	Hg
264	Tl	264	Tl
265	Pb	265	Pb
266	Bi	266	Bi
267	Po	267	Po
268	At	268	At
269	Hg	269	Hg
270	Tl	270	Tl
271	Pb	271	Pb
272	Bi	272	Bi
273	Po	273	Po
274	At	274	At
275	Hg	275	Hg
276	Tl	276	Tl
277	Pb	277	Pb
278	Bi	278	Bi
279	Po	279	Po
280	At	280	At
281	Hg	281	Hg
282	Tl	282	Tl
283	Pb	283	Pb
284	Bi	284	Bi
285	Po	285	Po
286	At	286	At
287	Hg	287	Hg
288	Tl	288	Tl
289	Pb	289	Pb
290	Bi	290	Bi
291	Po	291	Po
292	At	292	At
293	Hg	293	Hg
294	Tl	294	Tl
295	Pb	295	Pb
296	Bi	296	Bi
297	Po	297	Po
298	At	298	At
299	Hg	299	Hg
300	Tl	300	Tl
301	Pb	301	Pb
302	Bi	302	Bi
303	Po	303	Po
304	At	304	At
305	Hg	305	Hg
306	Tl	306	Tl
307	Pb	307	Pb
308	Bi	308	Bi
309	Po	309	Po
310	At	310	At
311	Hg	311	Hg
312	Tl	312	Tl
313	Pb	313	Pb
314	Bi	314	Bi
315	Po	315	Po
316	At	316	At
317	Hg	317	Hg
318	Tl	318	Tl
319			

[illegible]

Diphenylcarbazone as mercurimetric Indicator. De-

b. **Mermination of thiocyanates and of cyanides.** J. Tittlek, *Collection Czechoslov. Chem. Commun.* 10, 242 R (1938); cf. C. A. 32, 12057.—If a soln. of alkali or alk. earth thiocyanate in 3–60 ml. of water and 0.5–5 ml. of 0.2 N HNO_3 is treated with 1–3 drops of a cold, satd. soln. of diphenylcarbazone in alc., then, on titrating the mixt. with 0.1 N $Hg(NO_3)_2$, the end point will be reached when the white ppt. assumes a blue-violet tint, or, if the soln. is very dil., when the soln. assumes a rose-violet color. The results are usually within 0.4% of the truth. In the presence of heavy metal ions the results are somewhat different. When Fe^{+++} is present, $Fe(CNS)_3$ is formed which is its own indicator. Cu^{++} reacts with thiocyanate and the ppt. fixes the indicator so that it no longer reacts with $Hg(NO_3)_2$. If green Cr^{+++} is present a dark red thiocyanate is formed which does not change on adding the diphenylcarbazone but toward the end of the titration, when the thiocyanate attached to the Cr comes into reaction with the Hg , the color of the Cr returns and it is very easy to recognize the blue tint. For the detn. of cyanide 3 methods of carrying out the analysis were tested. (1) The soln. of cyanide was treated with an excess of $Hg(NO_3)_2$, made acid with 2 ml. of 0.2 N HNO_3 and then an excess of chloride or thiocyanate was added and this last excess titrated with $Hg(NO_3)_2$. (2) The excess of

A 18-51 A METALLURGICAL LITERATURE CLASSIFICATION

7

Micro-volumetric analysis with diphenylcarbohydrazide and diphenylcarbohydrazone as indicators (mercurimetry). J. V. LUKASZ AND J. LUTIKHA. *Microchem. J.* 11: 29 (1968) - Mercuric ions react with diphenylcarbohydrazide or with the corresponding hydrazone, to give an intensive violet-blue coloration so that the end point of the reaction between Hg^{++} and Cl^- to form undissocd. $HgCl_2$ can be detd. accurately when one of these org. substances is present as indicator in a soln. Procedures are given for standardizing $Hg(NO_3)_2$ solns. and titrating Cl^- in solns. of KCl , $BaCl_2$ and $CuCl_2$. The results obtained were excellent.

W. T. H.

ASB. S.L.A. METALLURGICAL LITERATURE CLASSIFICATION

7

Mercurimetric determination of iodine with diphenylcarbazone as indicator
 J. V. Dubsky and J. Fritlek. Chem. Zvest 8, 41-2 (1962) in English. Diphényl-
 carbazone can be used as a very sensitive indicator in the mercurimetric detn. of I.
 The intensively violet coloration with Hg^{2+} is clearly visible even in the presence of
 insol. HgI_2 formed during the reaction.
 Jaroslav Kubyra

ASB-51A METALLURGICAL LITERATURE CLASSIFICATION

62-2-1

11

CA

PROCESSES AND PROPERTIES INDEX


Diphenylcarbazide as indicator in mercurimetry. J. TRJILEK. Chem. Abstr. B, 3-545 English (1933). Chlorides even in the presence of salts of heavy metals can be detected by titration with $Hg(NO_3)_2$ soln., with diphenylcarbazide or diphenylcarbazone as indicator. cf. C. A. 27, 2107.

JANOS AV. KUCERA

CLASSIC ELEMENT

ASAC 31 A METALLURGICAL LITERATURE CLASSIFICATION

11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100



A-3

BZ

Oxidation of ethyl carbonate. J. V. DUMAY
and J. TAVLAK (Coll. Czech. Chem. Comm., 1983, 8,
49—60; cf. A., 1990, 199).—Oxidation of Et carb-
amate with I, H₂O₂ (acid or neutral), or CuCl₂ gives
3 : 5-dioctoxy-1 : 2 : 4-thiodiazole and S. A. A. L.

MATERIALS INDEX

ASB-SLA METALLURGICAL LITERATURE CLASSIFICATION

SUBJECT INDEX ONLY

COLLECTION

cc

Diphenylcarbazone as a mercurimetric indicator in the determination of bromine. J. Titlck. Chem. Abstr. 12, 1876 (1937).—Sols. of KBr or KCl were mixed with 0.1 cc. of a 1% soln. of diphenylcarbazone in KIOLH, treated with several drops of 0.2 N HNO₃, and titrated with 0.1 N Hg(NO₃)₂ in the cold. The first excess of Hg gave a violet color with diphenylcarbazone which indicator is specific for Hg ions. The best results were obtained when the soln. was 0.2 N in HNO₃. The presence of Cu, Fe, Pb, Zn and Mn did not affect the accuracy of the Br detns. if the solns. were 0.1 N in acid. Microtitrations with 0.002 N Hg(NO₃)₂ were satisfactory up to a concn. of 0.08 mg. Br per cc.; in more dil. solns. the presence of other metals and the HNO₃ affected the end point and gave low results.

Frank Marsh

118

Determination of magnesium in biological materials
An oxidation method. John P. Nielsen. *Ind. Eng. Chem., Anal. Ed.* 11, 610-611 (1939). Cf. C. A. 26, 4351; *Chem. Abstr.* 1939, 2. Mg pptd. with 8-hydroxyquinoline is detd. by oxidation with ammonium hexanitratocerate and titration of excess cerate with ferrous ammonium sulfate; o-phenanthroline ferrous sulfate is used as indicator. Analyses of canned tomato ash by this method agree with those by A. O. A. C. tentative method. J. McAfee.

The determination of small quantities of iodine in biological materials. L. Trillick. *Chem. Abstr.* 14, 1968 (1939).—In a modified Leichert app. (C. A. 27, 4266) without any rubber tubing, with a dephlegmator between the distg. flask and condenser, and without the useless 2nd condenser vessel, I. oxidized the org. substances with CrO_3 in the presence of H_2SO_4 and traces of CeSO_4 , reduced the iodic acid with Na_2SO_3 and volatilized the free I under low pressure into NaOH . After the soln. was concd., the I was oxidized with H_2O_2 and titrated by the Winkler method. The modified method eliminated all of the criticisms raised against the original app. Adds of 1-5 g of I to 10 cc. of blood were detd. with an accuracy of 10-20%. In 50-cc. urine samples, which could be concd., the accuracy of the I detn. was higher than 10%. Frank Marsh.

10

la

Oxidation of xanthamide. J. V. DUBSKÝ AND J. TETLICK. *Chem. Abstr.* 8, 1-2 (in English 3)(1933). The red coloration of xanthamide (I) with CuCl_2 can be explained by intermediate formation of compds. of the general formula $\text{CuCl}_2(\text{HCl})$, followed by the final sepn. of the addn. compd. CuCl_2 . Simultaneously there is formed the compd. $\text{C}_4\text{H}_5\text{N}_3\text{O}_2\text{S}$ which can be also prepd. by the oxidation of xanthamide with H_2O_2 or alk. I in HCl or neutral soln. The constitution of this compd. is explained by the formation of a thio-1,3-diazole $\text{N}:\text{C}(\text{OEt})\text{N}:\text{C}(\text{OEt})\text{S}$, m. 49-50°.

JAROMÍR KUČERA

ASR-5LA METALLURGICAL LITERATURE CLASSIFICATION

10

CR

Oxidation of xanthamide (contribution to the explanation of the reaction of this amide with cupric chloride). J. V. DUBSKÝ AND J. TRTILSKÝ. *Collection Czechoslov. Chem. Communications* 5, 40 60(1933). - See C. A. 27, 2137. ALFRED HOFFMANN

ASAC-SLA METALLURGICAL LITERATURE CLASSIFICATION

SECTION	SUBSECTION	CLASSIFICATION
1	1	1
2	2	2
3	3	3
4	4	4
5	5	5
6	6	6
7	7	7
8	8	8
9	9	9
10	10	10
11	11	11
12	12	12
13	13	13
14	14	14
15	15	15
16	16	16
17	17	17
18	18	18
19	19	19
20	20	20
21	21	21
22	22	22
23	23	23
24	24	24
25	25	25
26	26	26
27	27	27
28	28	28
29	29	29
30	30	30
31	31	31
32	32	32
33	33	33
34	34	34
35	35	35
36	36	36
37	37	37
38	38	38
39	39	39
40	40	40
41	41	41
42	42	42
43	43	43
44	44	44
45	45	45
46	46	46
47	47	47
48	48	48
49	49	49
50	50	50
51	51	51
52	52	52
53	53	53
54	54	54
55	55	55
56	56	56
57	57	57
58	58	58
59	59	59
60	60	60
61	61	61
62	62	62
63	63	63
64	64	64
65	65	65
66	66	66
67	67	67
68	68	68
69	69	69
70	70	70
71	71	71
72	72	72
73	73	73
74	74	74
75	75	75
76	76	76
77	77	77
78	78	78
79	79	79
80	80	80
81	81	81
82	82	82
83	83	83
84	84	84
85	85	85
86	86	86
87	87	87
88	88	88
89	89	89
90	90	90
91	91	91
92	92	92
93	93	93
94	94	94
95	95	95
96	96	96
97	97	97
98	98	98
99	99	99
100	100	100

10

Ca

Hippurhydroxamic acid. J. V. DUBAKÝ AND J. TRTILSKÝ. *Collection Czechoslov. Chem. Communications* 3, 103-11(1933).—Hippuramide, white needles, m. 181°, treated in aq. alc. with 2 mols. $\text{NH}_4\text{OH} \cdot \text{HCl}$ and NaOAc yields, not the amidoxime, but the free hydroxamic acid (I) in 2 forms: *cis*, white scales, m. 141°, decompd. on drying at 100°, prepd. below 70°, and *trans*, needles, m. 159°, prepd. at 100°, each sol. in EtOH and hot H_2O . Both forms of I with $\text{Cu}(\text{OAc})_2$ give a quant. ppt. of the green $\text{BzNHCH}_2\text{C}(\text{NO})\text{Cu}_2\text{O}$ sol. in acid; with CoCl_2 complex mixts. contg. Co^{++} are obtained. $\text{Hg}(\text{NO}_3)_2$ and I yield the yellow, unstable $\text{BzNHCH}_2\text{C}(\text{NOH})_2 \cdot 3 \text{H}_2\text{O}$. FeCl_3 gives the intensely red $(\text{BzNHCH}_2\text{C}(\text{OH})\text{NO})_2\text{FeOH} \cdot \text{H}_2\text{O}$. H. A. B

458-35A METALLURGICAL LITERATURE CLASSIFICATION

112

ca

Determination of chlorides in urine and blood with diphenylcarbazone as indicator. J. Krulik. *Bratslavskii Lekarski List* 17, No. 10, 295 (1937). Reagents: (1) 0.1 N soln. $Hg(NO_3)_2$ (to 10.23 g. of $Hg(NO_3)_2$ is added 2.4 cc. of concd. HNO_3 , the contents are brought to 1 l., filtered and the soln. is standardized against pure $NaCl$); (2) a satd. soln. of diphenylcarbazone in alk. in $NaCl$; (3) 2 N HNO_3 , free from Cl . To 2.3 cc. of the cold, (4) 2 N HNO_3 and dild. to 100 urine, acidified with 1.3 cc. of 2 N HNO_3 and dild. to 100 (50 cc. is added 0.5 cc. of diphenylcarbazone. The contents are titrated with $Hg(NO_3)_2$ to a violet-rose color and the amt. of Cl calcd. One cc. of blood, in a test tube, having been acidified with 2 cc. of 2 N HNO_3 , heated to boiling, dild. to 10 cc. and again heated to boiling, is dild. to 50 cc. in a volumetric flask and filtered. The clear filtrate is titrated with 0.01 N $Hg(NO_3)_2$ from a microburet with diphenylcarbazone as the indicator.

V. D. Karpenko

ASB-51A METALLURGICAL LITERATURE CLASSIFICATION

12-

PROCESSING AND PROPERTIES INDEX

Diphenylcarbazone as an indicator for mercury for the determination of bromides. J. TRILLER (Chem. Abstr. 1937, 12, 184-185).--The highest acidity at which the mercurimetric determination of Br⁻ using diphenylcarbazone as an indicator can be carried out (even in presence of ions of heavy metals) is 0.2N-HNO₃. The best condition is neutral or very slightly acid. F. R.

ASD-SLA METALLURGICAL LITERATURE CLASSIFICATION


SECTION	SUBSECTION	NUMBER	DATE	REMARKS
1	1	1	1	1
2	2	2	2	2
3	3	3	3	3
4	4	4	4	4
5	5	5	5	5
6	6	6	6	6
7	7	7	7	7
8	8	8	8	8
9	9	9	9	9
10	10	10	10	10
11	11	11	11	11
12	12	12	12	12
13	13	13	13	13
14	14	14	14	14
15	15	15	15	15
16	16	16	16	16
17	17	17	17	17
18	18	18	18	18
19	19	19	19	19
20	20	20	20	20
21	21	21	21	21
22	22	22	22	22
23	23	23	23	23
24	24	24	24	24
25	25	25	25	25
26	26	26	26	26
27	27	27	27	27
28	28	28	28	28
29	29	29	29	29
30	30	30	30	30
31	31	31	31	31
32	32	32	32	32
33	33	33	33	33
34	34	34	34	34
35	35	35	35	35
36	36	36	36	36
37	37	37	37	37
38	38	38	38	38
39	39	39	39	39
40	40	40	40	40
41	41	41	41	41
42	42	42	42	42
43	43	43	43	43
44	44	44	44	44
45	45	45	45	45
46	46	46	46	46
47	47	47	47	47
48	48	48	48	48
49	49	49	49	49
50	50	50	50	50
51	51	51	51	51
52	52	52	52	52
53	53	53	53	53
54	54	54	54	54
55	55	55	55	55
56	56	56	56	56
57	57	57	57	57
58	58	58	58	58
59	59	59	59	59
60	60	60	60	60
61	61	61	61	61
62	62	62	62	62
63	63	63	63	63
64	64	64	64	64
65	65	65	65	65
66	66	66	66	66
67	67	67	67	67
68	68	68	68	68
69	69	69	69	69
70	70	70	70	70
71	71	71	71	71
72	72	72	72	72
73	73	73	73	73
74	74	74	74	74
75	75	75	75	75
76	76	76	76	76
77	77	77	77	77
78	78	78	78	78
79	79	79	79	79
80	80	80	80	80
81	81	81	81	81
82	82	82	82	82
83	83	83	83	83
84	84	84	84	84
85	85	85	85	85
86	86	86	86	86
87	87	87	87	87
88	88	88	88	88
89	89	89	89	89
90	90	90	90	90
91	91	91	91	91
92	92	92	92	92
93	93	93	93	93
94	94	94	94	94
95	95	95	95	95
96	96	96	96	96
97	97	97	97	97
98	98	98	98	98
99	99	99	99	99
100	100	100	100	100

137 AND 138 SERIES										139 AND 140 SERIES									
PROCEDURES AND PROPERTIES INDEX																			
<p><i>EC</i></p> <p><i>A-1</i></p> <p>(1) Microchemical analysis of the following compounds: ortho- and para-phenylenediamine, and ortho- and para-aminophenol. J. V. Deane and E. J. Deane, <i>Microchem. J.</i> 1966, 12: 222-223. The authors give an intense violet-blue coloration with ortho-phenylenediamine (H) and ortho-aminophenol (H). The color may be titrated in 0.1M sodium acetate with 0.01N $Hg(NO_3)_2$ using (H) as buffer (H). In the case of ortho-aminophenol, the formation of non-colored $HgCl_2$ by metals have little disturbing effect; in presence of the results were 0.5% low.</p> <p>J. S. A.</p>																			
MATERIALS INDEX										COMMON VARIANTS INDEX									
ASR-11A METALLURGICAL LITERATURE CLASSIFICATION										FROM SOURCE									
FROM SYNDICATE										FROM SOURCE									
100000 01										011111 011111 111									
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20										1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20									

The color reaction of bismuth. II. J. V. Dubsky and
 J. Trtilka, *Chem. Obsor* 9, 243-5 (in English 245) (1934).
 The K salt of mercaptophenylthiothiazolone is just as
 sensitive a reagent for Bi as dimercaptothiothiazole (bis-
 muthiol I). The limit of detection is 1.2 γ Bi; the limit of
 diln. 1:28,000. J. Kucera

Organic reagents in analytical chemistry. IV. J

Dubinsky and J. Lukac¹, *Chem. Abstr.* 9, 112,4 (1915), cf. C. A. 20, 5017. For the following reagents are reviewed: 2,5-dimercapto-1,3,4-thiadiazole, 5-mercapto-3-phenyl-2-thio-1,3,4-thiadiazole-2-one, cantharidin, by dioxysquindones, pyrogallol, $K_2Cr_2O_7$ -Na₂ tartrichloride, cobalt chloride, thionine, bromine, dimethyl oxamine, nitrophenylhydrosulfoxamine, thalamine B, glutamine, nitrophenylhydrosulfoxamine, thalamine B, thiophosphorylcarbamate, antipyrine, piperazine, $[Co(CN)_5]^-$, K_2 , quinine, phenylthiohydantonic acid, viscose, urotopine and hematein. Other literature is also compiled up to date. J. Kucera



ca
The reaction of aminobenzoic acids with cadmium and zinc salts. J. V. Trifilek. *Chem. Listy* 29, 70 (1935). A soln. of the free aminobenzoic acids did not react with either Zn or Cd salts; after an addn. of Na acetate to the free acids, only the *o*-acid formed a ppt. with Cd and Zn. In a neutral or faintly acid soln., *o*-NH₂C₆H₄COOK gave a white ppt. immediately with Zn (even in 0.1 N soln.) but gave a white ppt. with Cd after many min.; *p*-NH₂C₆H₄COOK (I) did not react visibly with Zn but yielded an instantaneous, white, cryst. ppt. with Cd. While the Zn salts of the aminobenzoic acids were very insol., those of Cd were very insol. in a range of conditions as in a precisely neutral medium, in a soln. of the *p*-acid neutralized with KOH against phenol-

phthalein, and in the presence of an excess of basic salt of the aminobenzoic acid. The Cd ppt. with *p*-NH₂C₆H₄COOK (CdHCOO(NH₂C₆H₄), 2H₂O) was only slightly sol. in free H₂O and 50% EtOH, dissolved quickly in weak acids or in weakly acidified soln., and could not be used for quant. analysis. With 1 cc. of 0.01 N CdSO₄, I gave a ppt. instantly; with 1 cc. of 0.001 N CdSO₄, it gave a ppt. after 30 min.; this corresponded to 0.55 mg. of Cd in a threshold concn. of 1.1780. Under identical conditions, ZnSO₄ did not yield a ppt. (only the 1.0 N Zn soln. gave a temporary ppt. which quickly dissolved and reappeared only in an excess of the Zn reagent as a white lake). The reactions of the *o*, *m*, and *p*-aminobenzoic acids with neutral soln. with Ag, Pb, Hg, Cu, Sn, Bi, Sb, Co, Ni and Fe are described; all ppts. dissolved in dil. HNO₃.
Frank Maresh

The reaction of bismuth salts with the condensation products of bisphenylthioureas. J. A. Dinkels and J. E. Little. (*Ann. Inst. 20, 31 4-1935*). In EtOH, 0.005 g. mol. BiCl₃ with 0.015 mol. 3-amino-5-mercapto-4-phenyl-1,2,4-triazole gave a yellow ppt. which dissolved to form a yellow soln. from which a fine orange ppt. settled. After the soln. was reheated, the orange ppt. redissolved and left a mucous red mass in the tube. The yellow ppt. could not be produced from the soln. again. In concd. soln. more of the orange than of the yellow ppt. formed. Pb and Sb formed a yellow white ppt., Ag, Hg, Cd, Zn and Ba, a white one; Cu, a blue-black one, Co, a pink one and Ni (after an addn. of Na acetate), a green-white ppt. In EtOH, BiCl₃ with 5-mercapto-3-amino-4-thio-1,2-diazole gave a yellow ppt. This reaction is analogous to those of thioureas. In dil. EtOH solns. BiCl₃ did not react with 3,5-di-amino-4-thio-1,2-diazole, in N soln. it formed a white ppt.; Ag, Hg⁺⁺, and Hg⁺ ppts. were yellow-white; Pb, Sb and Cd were white; Cu was brown-black, Ni was green, Co was rose, Fe was brick red. In EtOH, BiCl₃ with 3,5-dimercapto-4-phenyl-1,2,4-triazole formed an orange ppt., BiCl₃·2N₂S₂·CHCl₃, in the presence of an excess of BiCl₃ the ppt. was red-orange. Pb and Sb yielded yellow salts; Ag yielded a yellow-white ppt.; Co formed an olive-green ppt. The Bi reaction in the last case is analogous to that of the Bi thiole. Frank March

Frank Marshall

CIA-RDP86-00513R001756810008-5"

Calcium hetero formate compounds. J. V. Dubsky
and J. Trtielek. *Pub. facult. sci. univ. Masaryk No. 100*,
3-6 (1934). From $[\text{Ca}_2(\text{form})_2\text{Cl}_2 \cdot 8\text{H}_2\text{O}]$, in which "form"
is HCOO^- , the following complex compds. were prepd.:
 $[\text{Ca}_2(\text{form})_2][\text{O}(\text{NO}_2)_2\text{C}_6\text{H}_3] \cdot 4\text{H}_2\text{O}$ (picrate), $[\text{Ca}_2(\text{form})_2] \cdot 3$
 $\text{CrO}_4 \cdot 2\text{H}_2\text{O}$, $[\text{Ca}(\text{form})_2] \cdot \text{K}_2\text{Fe}(\text{CN})_6$ and $[\text{Ca}_2(\text{OAc})_2] \cdot$
form, or $[\text{Ca}_2(\text{form})_2(\text{OAc})_2]$. V. D. Karpenko

ASH S.L.A. METALLURGICAL LITERATURE CLASSIFICATION

The formation of salts of amidoximes I Hippuramidoxime. J. V. Dulack and J. H. Dole, *Collection Czechoslov. Chem. Communications*, 8, 100 (1943). Hippuramidoxime (I), m. 123-6°, was prepd. from hippuronitrile and H_2NOH , but could not be obtained free from Cl $_2$. It gave a reddish violet color with CuCl_2 and a brownish green ppt. with $\text{Cu}(\text{OAc})_2$. This Cu salt, with a ratio to Cu:N of 1:3, contained 2 mol% of H_2O , only one of which could be driven off below 100°; so the authors gave it the formula: $\text{C}_9\text{H}_{11}\text{CONHC}(=\text{N})(\text{NH}_2)\cdot\text{NOCuOH}_2\cdot\text{H}_2\text{O}$, and for the formula of I: $\text{C}_9\text{H}_{11}\text{CONHC}(=\text{N})(\text{NH}_2)\cdot\text{NOH}$. NiCl_2 alone has no action on I but the addn. of a little NH_4OH causes a very slow formation of a red-violet ppt. This action is greatly accelerated by a little H_2O_2 . This ppt. appears to be a mixt. of 2 nickelic salts: $\{\text{C}_9\text{H}_{11}\text{CONHC}(=\text{N})(\text{NH}_2)\cdot\text{NO}\}_2\cdot\text{NiOH}$ and $\{\text{C}_9\text{H}_{11}\text{CONHC}(=\text{N})(\text{NH}_2)\cdot\text{NO}\}_2\cdot\text{Ni}$.

John E. Millery

John E. Millery

4.1.3.1 METALLURGICAL LITERATURE CLASSIFICATION

CA

The formation of salts of amidoximes I Hippur
amidoxime J. V. Dubsky and J. Fritsch Collection
Czechoslov Chem Communications 8, 1011 (1963).
Hippuramidoxime (I), m. 123.0°, was prepd from
hippuramitrite and H_2NOH , but could not be obtained
free from Cl_2 . It gave a reddish violet color with aq
 $FeCl_3$ and a brownish green ppt with $CoCl_2 \cdot 6H_2O$. This
 Co salt, with a ratio to Co N of 1.8, contained 2 mols
of H_2O , only one of which could be driven off below 100°
so the authors gave it the formula $C_9H_{11}CONHCH_2$
 $C(NH_2) \cdot NO \cdot Co(OH) \cdot 2H_2O$, and for the formula of I: C_9H_{11}
 $CONHCH_2C(NH_2) \cdot NOH$. $NiCl_2$ alone has no action
on I but the addn. of a little NH_4OH causes a very slow
formation of a red-violet ppt. This action is greatly
accelerated by a little H_2O_2 . This ppt appears to be a
mixture of 2 nickelous salts: $[C_9H_{11}CONHCH_2C(NH_2) \cdot NO]_2 \cdot$
 NOH and $[C_9H_{11}CONHCH_2C(NH_2) \cdot NO]_2 \cdot Ni$.

John E. Millery

ASH-35A METALLURGICAL LITERATURE CLASSIFICATION

Acetato compounds of the alkaline earth metals. J. V. DUNN AND J. TRIFER.
Pub. Chem. Soc. New York, 1932, No. 161, 1-29 (in English, 30). Cf. Weinland and
Hachenberg, C. A. 17, 2841. By a series of direct syntheses and a series of substitutions
of the free anion groups the following acetato compounds were prepared: $\text{Ba}(\text{C}_2\text{H}_3\text{O}_2)_2$,
 $[\text{Ca}_{\text{ac}}]_2[\text{Cr}, 10\text{H}_2\text{O}]$, $[\text{Ca}_{\text{ac}}][\text{Sr}, 10\text{H}_2\text{O}]$, $[\text{Ca}_{\text{ac}}][\text{Ba}, 10\text{H}_2\text{O}]$, $[\text{Ca}_{\text{ac}}](\text{NO}_3)(\text{CH}_3\text{COOH})$,
 $[\text{Ca}_{\text{ac}}](\text{per}, 6\text{H}_2\text{O})$, $[\text{Ca}_{\text{ac}}][\text{CrO}_2, 2\text{H}_2\text{O}]$, $[\text{Ca}_{\text{ac}}][\text{CrO}_2, \text{CH}_3\text{COOH}]$, $[\text{Ca}_{\text{ac}}][\text{acSCN}, 3\text{H}_2\text{O}]$,
 $[\text{Ca}_{\text{ac}}](\text{SO}_4, 4\text{H}_2\text{O})$, $[\text{Ca}_{\text{ac}}](\text{NO}_3), \text{Ca}(\text{ac}), \text{Sr}(\text{ac}), [\text{Sr}_{\text{ac}}](\text{Cl}, 2\text{H}_2\text{O})$, $[\text{Sr}_{\text{ac}}](\text{Cl}, \text{Sr},$
 $\text{Cl}, 14\text{H}_2\text{O})$, $[\text{Sr}_{\text{ac}}][\text{acCl}, 7\text{H}_2\text{O}]$, $[\text{Mg}_{\text{ac}}][\text{acCl}, 5\text{H}_2\text{O}], 6\text{H}_2\text{O}$. The assumptions con-
cerning the constitution of these compds. are based on analogy with the complex acetato
compds. of Ba (cf. Weinland and Henrichsen, C. A. 17, 1509) and of Pb (cf. Weinland and
Paul, C. A. 17, 3263), on numerous substitution reactions, and on such properties of the
compds. themselves as water of crystal.

M. G. MOORE

M. G. Mendenhall

ASR 514 METALLURGICAL LITERATURE CLASSIFICATION

[illegible]

BC

A 3

Note: In preparation of the products of condensation of 1-phenylthiocarbamide, J. V. Duncanson and J. T. H. (Chem. Lett., 1935, 20, 33-34).— HCl , yields an orange-yellow salt with 3-anilino-5-thiol-4-phenyl-1:2:4-triazole; an orange-red salt, $\text{HC}_6\text{H}_4\text{N}_3\text{S}_2\text{Cl}_2 \cdot \text{H}_2\text{O}$, with 3:5-dithiadiphenyl-1:2:4-triazole. A white salt with 3:5-dithiadiphenyl-1:2:4-triazole and a yellow salt with 3-anilino-5-thio-4-phenyl-1:2:4-triazole. R. T.

ASM-SLA METALLURGICAL LITERATURE CLASSIFICATION

1ST AND 2ND DEGREE										3RD AND 4TH DEGREE									
PROCESSES AND PROPERTIES INDEX																			
BC										A-1									
<p>DUSKÝ, J. V., and TRTILK, J. A contribution to the study of acetate-compounds of the alkaline-earth metals. (<i>Publ. Fac. Sci. Univ. Masaryk</i>, No. 181). Brno 1933. pp. 30. [In Czech. English summary.]</p>																			
ASB-ELA METALLURGICAL LITERATURE CLASSIFICATION																			
FROM STUDYING										FROM WORKING									
100000 110 120 130 140 150 160 170 180 190 200										210 220 230 240 250 260 270 280 290 300									
C																			

TRTILEK, Josef

Chemie pro osmy postupny rocnik vseobecne vzdelavacich skol. (Chemistry for the 8th Grade of the Schools of General Education. 4th ed. illus., index) Prague, SPN, 1957. 143 p.

Bibliograficky katalog, CSR, Ceske knihy, No. 33. 24 Sept 57. p. 712.

5.111-8, J.

J. V. DUSIN, Mikrodiam, 1985, 17, 323-343

TRTILEK, J.

J. V. DUBSKY, Publ. Fac. Sci. Univ. Masaryk, 1936, 223, 7-10

TRTILEK, J.,
J. V. LUBSKY, (Chem. Obzor, 1934, 9, 173-174)

TUTILEK, J.

J. V. DUMKY, Coll. Czech. Ther. Commun. 1, 141-3, 1936

TCHIEV, A.

A. A. TCHIEV, Pub. Faculte sci. univ. Masaryk No. 223, 7-10, 1936

TRTILEK, J.

J. V. DUBSKY, Collection Czechoslov. Chem. Communications 7, 1-9, 1935